

Factor XIIIa Rabbit mAb

货号: **AYM30634**

产品信息

| | |
|-------|--|
| 反应 | Human |
| 宿主 | Rabbit |
| 克隆性 | Monoclonal |
| 预测反应 | |
| 应用 | WB IHC IF/ICC FC |
| 推荐浓度 | WB: 1:500 - 1:2000 IHC: 1:50 - 1:200 IF/ICC: 1:50 - 1:200 FC: 1:20 - 1:50 |
| 理论分子量 | 83kDa |
| 实测分子量 | 83kDa |
| 形式 | Liquid |
| 保存条件 | Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.75% BSA,50% glycerol,pH7.3. |
| 偶联物 | Unconjugated |
| 阳性对照 | Jurkat,SH-SY5Y,BxPC-3,Mouse testis,Rat brain,Rat testis |
| 细胞定位 | Cytoplasm,Secreted |
| 纯化 | Affinity purification |


抗原信息

| | |
|------|--|
| 抗原信息 | Recombinant fusion protein corresponding to Human Factor XIIIa. |
| 序列 | LEQASLHFFVTARINETRDVLAKQKSTVLTPIEIIKVRGTQVVGSDMTVTVEFTNPLKETLRNVVWHLDGPVTRPMKKMF REIRPNSTVQWEEVCRPWVSGHRKLIASMSDSL RHVYGELDVQIQRPSM |

靶点信息

| | |
|-------|---|
| 研究背景 | This gene encodes the coagulation factor XIII A subunit. Coagulation factor XIII is the last zymogen to become activated in the blood coagulation cascade. Plasma factor XIII is a heterotetramer composed of 2 A subunits and 2 B subunits. The A subunits have catalytic function, and the B subunits do not have enzymatic activity and may serve as plasma carrier molecules. Platelet factor XIII is comprised only of 2 A subunits, which are identical to those of plasma origin. Upon cleavage of the activation peptide by thrombin and in the presence of calcium ion, the plasma factor XIII dissociates its B subunits and yields the same active enzyme, factor XIIIa, as platelet factor XIII. This enzyme acts as a transglutaminase to catalyze the formation of gamma-glutamyl-epsilon-lysine crosslinking between fibrin molecules, thus stabilizing the fibrin clot. It also crosslinks alpha-2-plasmin inhibitor, or fibronectin, to the alpha chains of fibrin. Factor XIII deficiency is classified into two categories: type I deficiency, characterized by the lack of both the A and B subunits; and type II deficiency, characterized by the lack of the A subunit alone. These defects can result in a lifelong bleeding tendency, defective wound healing, and habitual abortion. |
| 基因ID | 2162 |
| 基因名 | F13A1 |
| Swiss | P00488 |
| 别名 | F13A1;F13A |

产品验证



Western blot analysis of Factor XIIIa expressed in Jurkat,SH-SY5Y,BxPC-3 using Factor XIIIa Rabbit mAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn