

NAT10 Rabbit mAb

货号: **AYM30636**

产品信息

反应	Human,Mouse,Rat
宿主	Rabbit
克隆性	Monoclonal
预测反应	
应用	WB IHC IF/ICC IP FC
推荐浓度	WB: 1:500 - 1:2000 IHC: 1:50 - 1:200 IF/ICC: 1:50 - 1:200 IP: 1:20 - 1:50 FC: 1:20 - 1:50
理论分子量	116kDa
实测分子量	116kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.75% BSA,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	HeLa,K-562,SH-SY5Y,Mouse testis
细胞定位	Nucleus,nucleolus
纯化	Affinity purification

抗原信息


抗原信息	Recombinant fusion protein corresponding to Human NAT10.
序列	KTLTDEDEADQGGWLAAFWKDFRRRFLALLSYQFSTFSPSLALNIIQNRNMGKPAQPALSREELEALFLPYDLKRLEMYSR NMVDYHLIMDMIPASRIYFLNQLGDLALSAAQSALLLIGLQHKSVQLEKEIELPSGQLMGLFNRIIRKVVKLFNEVQEKA EEQMVAAKDVMEPTMKTLSDDLDEAAKEFQEKHKKEVGLKSMDSLSEYIIRGDDEEWNEVLNKAGPNASIIISLKSDDK KLEAKQEPKQSKLKNRETKNKKDMKLRKK

靶点信息

研究背景	The protein encoded by this gene is an RNA cytidine acetyltransferase involved in histone acetylation, tRNA acetylation, the biosynthesis of 18S rRNA, and the enhancement of nuclear architecture and chromatin organization.
------	--

基因ID	55226
基因名	NAT10
Swiss	Q9H0A0
别名	NAT10; ALP; NET43; Kre33; N-acetyltransferase 10

产品验证



Western blot analysis of NAT10 expressed in Mouse testis, HeLa, K-562 using NAT10 Rabbit mAb at 1:10 00. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn