

QK1 Rabbit mAb

货号: **AYM30643**

产品信息

| | |
|-------|---|
| 反应 | Human,Mouse |
| 宿主 | Rabbit |
| 克隆性 | Monoclonal |
| 预测反应 | |
| 应用 | WB IF/ICC IP |
| 推荐浓度 | WB: 1:500 - 1:2000 IF/ICC: 1:50 - 1:200 IP: 1:20 - 1:50 |
| 理论分子量 | 35kDa/36kDa/37kDa |
| 实测分子量 | 38kDa |
| 形式 | Liquid |
| 保存条件 | Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.75% BSA,50% glycerol,pH7.3. |
| 偶联物 | Unconjugated |
| 阳性对照 | HeLa,NIH/3T3,C2C12,C6 |
| 细胞定位 | Cytoplasm,Nucleus |
| 纯化 | Affinity purification |

抗原信息


| | |
|------|--|
| 抗原信息 | Recombinant fusion protein corresponding to Human QK1. |
| 序列 | MVGEMETKEKPKPTPDYLMQLMNDKMLMSSLPNFCGIFNHLERLLDEEISVRKDMYNDTLNGSTEKRSAELPDAVGPIV QLQEKLYVPVKEYPDFNFVGRILGPRGLTAKQLEAETGCKIMVRGKGSMRDKKKEEQNRGKPNWEHLNEDLHVLITVEDA QNRAEIKLKRAVEEVKLLVPAEGEDSLKMKQLMELAILNGTYRDANIKSPALAFSLAATAQAAPRIITGPAPVLPAAALRTP TPAGPTIMPLIRIQITAVMPNGTPHPTAAIVPPGPEAGLIYTPYEYPYTLAPATSILEYPIEPSGVLGAVATKVRRHDMRVHPYQ RIVTADRAATGN |

靶点信息

| | |
|------|--|
| 研究背景 | The protein encoded by this gene is an RNA-binding protein that regulates pre-mRNA splicing, export of mRNAs from the nucleus, protein translation, and mRNA stability. The encoded protein is involved in myelination and oligodendrocyte differentiation and may play a role in schizophrenia. Multiple transcript variants encoding different isoforms have been found for this gene. |
|------|--|

| | |
|-------|-------------------------|
| 基因ID | 9444 |
| 基因名 | QKI |
| Swiss | Q96PU8 |
| 别名 | QKI;Hqk;QK;QK1;QK3;hqkl |

产品验证



Western blot analysis of QKI expressed in HeLa, NIH/3T3, C2C12 using QKI Rabbit mAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn