

IRAK Rabbit mAb

货号: **AYM30646**

产品信息

反应	Human
宿主	Rabbit
克隆性	Monoclonal
预测反应	
应用	WB IF/ICC IP
推荐浓度	WB: 1:500 - 1:2000 IF/ICC: 1:50 - 1:200 IP: 1:20 - 1:50
理论分子量	68kDa/73kDa/74kDa/76kDa
实测分子量	77kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.75% BSA,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	HeLa,A-549,Mouse liver,Mouse kidney,Mouse skeletal muscle,Rat liver,Rat kidney,Rat skeletal muscle
细胞定位	Cytoplasm,Lipid droplet,Nucleus
纯化	Affinity purification

抗原信息


抗原信息	Recombinant fusion protein corresponding to Human IRAK.
序列	VKTHGARTKYLVIYERLEKIQAVVAGVPGHSEAASCIPPSPQENSYVSSTGRAHSGAAPWQPLAAPSGASAQAQAEQLQRG PNQPVESDESLGGLSAALRSWHLTPSCPLDPAPLREAGCPQGDGTAGESSWGS GPGSRPTAVEGLALGSSASSSEPPQII INPARQKMVQKLALYEDGALDSLQLLSSSSLPGLGLEQDRQGP EESDEFQS

靶点信息

研究背景	This gene encodes the interleukin-1 receptor-associated kinase 1, one of two putative serine/threonine kinases that become associated with the interleukin-1 receptor (IL1R) upon stimulation. This gene is partially responsible for IL1-induced upregulation of the transcription factor NF-kappa B. Alternatively spliced transcript variants encoding different isoforms have been found for this gene.
------	---

基因ID	3654
基因名	IRAK1
Swiss	P51617
别名	IRAK1;IRAK;pelle

产品验证



Western blot analysis of IRAK expressed in HeLa,A-549,Mouse liver using IRAK Rabbit mAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn