

Nucleostemin Rabbit mAb

货号: **AYM30654**

产品信息

反应	Human
宿主	Rabbit
克隆性	Monoclonal
预测反应	
应用	WB IF/ICC FC
推荐浓度	WB: 1:500 - 1:2000 IF/ICC: 1:50 - 1:200 FC: 1:20 - 1:50
理论分子量	60kDa/61kDa
实测分子量	65kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.75% BSA,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	293T
细胞定位	Nucleus,nucleolus
纯化	Affinity purification

抗原信息


抗原信息	Recombinant fusion protein corresponding to Human Nucleostemin.
序列	MTCHKRYKIQKKVREHHRKLRKEAKKRGHKKPRKDPGVPNSAPFKEALLREAELRKQRLEELKQQQLDRQKELEKRRKLE TNPDIKPSNVEPMEKEFGLCKTENKAKSGKQNSKKLYCQELKKVIEASDVVLEVL DARDPLGCRCPQVEEAIVQSGQKKLV LILNKSDLVPKENLESWLNLYLKKELPTVVFRASKPKDKGKITKRVKAKKNAAPFRSEVCFGKEGLWKLGGFQETCS

靶点信息

研究背景	The protein encoded by this gene may interact with p53 and may be involved in tumorigenesis. The encoded protein also appears to be important for stem cell proliferation. This protein is found in both the nucleus and nucleolus. Three transcript variants encoding two different isoforms have been found for this gene.
------	--

基因ID	26354
基因名	GNL3
Swiss	Q9BVP2
别名	GNL3;C77032;E2IG3;NNP47;NS

产品验证



Western blot analysis of Nucleostemin expressed in 293T using Nucleostemin Rabbit mAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn