

FOXP1 Rabbit mAb

货号: **AYM30673**

产品信息

反应	Human,Mouse,Rat
宿主	Rabbit
克隆性	Monoclonal
预测反应	
应用	WB
推荐浓度	WB: 1:500 - 1:2000
理论分子量	12kDa/54kDa/65kDa/66kDa/74kDa/75kDa/77kDa
实测分子量	80,90kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.75% BSA,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	HCT116,U-87MG,MCF7,HepG2,Mouse spleen,Mouse embryo
细胞定位	Nucleus
纯化	Affinity purification

抗原信息


抗原信息	Recombinant fusion protein corresponding to Human FOXP1.
序列	SGNPSLIKMQSSHAYCTPLNAALQASMAENSIPLYTTASMGNPTLGNLASAIREELNGAMEHTNSNESDSSPGRSPMQA VHPVHVKEEPDPEEAEGPLSLVTTANHSPDFDHRDYEDPVEDME

靶点信息

研究背景	This gene belongs to subfamily P of the forkhead box (FOX) transcription factor family. Forkhead box transcription factors play important roles in the regulation of tissue- and cell type-specific gene transcription during both development and adulthood. Forkhead box P1 protein contains both DNA-binding- and protein-binding domains. This gene may act as a tumor suppressor as it is lost in several tumor types and maps to a chromosomal region (3p14.1) reported to contain a tumor suppressor gene(s). Alternative splicing results in multiple transcript variants encoding different isoforms.
基因ID	27086

基因名	FOXP1
Swiss	Q9H334
别名	FOXP1;12CC4;HSPC215;MFH;QRF1;hFKH1B

产品验证



Western blot analysis of FOXP1 expressed in Mouse spleen, HCT116, U-87MG using FOXP1 Rabbit mAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn