

# IRF2 Rabbit mAb

货号: **AYM30678**

## 产品信息

反应	Human,Mouse
宿主	Rabbit
克隆性	Monoclonal
预测反应	
应用	WB IHC IF/ICC
推荐浓度	<b>WB:</b> 1:500 - 1:2000 <b>IHC:</b> 1:50 - 1:200 <b>IF/ICC:</b> 1:50 - 1:200
理论分子量	39kDa
实测分子量	50kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.75% BSA,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	Jurkat,Mouse lung
细胞定位	Nucleus
纯化	Affinity purification

## 抗原信息


抗原信息	Recombinant fusion protein corresponding to Human IRF2.
序列	KKGKKPKTEKEDKVKHIKQEPVSSLGSLNGVSDLSPEYAVLTSTIKNEVDSTVNIIVVGQSHLDSNIENQEIVTNPPDICQV VEVTTESDEQPVSMSSELYPLQISPVSSYAESSETTDSVPSDEESAEGRPHWRKRNIIEGKQYLSNMGTRGSYLLPGMASFVT SNKPDQLQVTIKESNPVYNSSWPPFQDLPLSSMTPASSSSRPDRETRASVIKKTSDITQARVKSC

## 靶点信息

研究背景	IRF2 encodes interferon regulatory factor 2, a member of the interferon regulatory transcription factor (IRF) family. IRF2 competitively inhibits the IRF1-mediated transcriptional activation of interferons alpha and beta, and presumably other genes that employ IRF1 for transcription activation. However, IRF2 also functions as a transcriptional activator of histone H4.
------	--

基因ID	3660
基因名	IRF2
Swiss	P14316
别名	IRF2;IRF-2

## 产品验证



Western blot analysis of IRF2 expressed in Jurkat, Mouse lung using IRF2 Rabbit mAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

## 实验步骤

访问官网浏览详情: [www.ablybio.cn](http://www.ablybio.cn)