

— ABLYBIO, Help Your Research



UAP1 Rabbit mAb

货号: **AYM30691**

产品信息

反应	Human,Mouse,Rat
宿主	Rabbit
克隆性	Monoclonal
预测反应	
应用	WB IHC IF/ICC IP FC
推荐浓度	WB: 1:500 - 1:2000 IHC: 1:50 - 1:200 IF/ICC: 1:50 - 1:200 IP: 1:20 - 1:50 FC: 1:20 - 1:50
理论分子量	59kDa
实测分子量	59kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.75% BSA,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	HepG2,DU145,Jurkat,Mouse brain,Rat brain
细胞定位	Cytoplasm
纯化	Affinity purification


抗原信息

抗原信息	Recombinant fusion protein corresponding to Human UAP1.
------	---

靶点信息

研究背景	Converts UTP and GlcNAc-1-P into UDP-GlcNAc, and UTP and GalNAc-1-P into UDP-GalNAc. Isoform AGX1 has 2 to 3 times higher activity towards GalNAc-1-P, while isoform AGX2 has 8 times more activity towards GlcNAc-1-P.
基因ID	6675
基因名	UAP1
Swiss	Q16222 (https://www.uniprot.org/uniprotkb/Q16222/entry)
别名	UAP1,UAP1 Rabbit mAb,Antigen X,Protein-pyrophosphorylation enzyme,Sperm-associated antigen 2,UDP-N-acetylgalactosamine pyrophosphorylase,UDP-N-acetylglucosamine pyrophosphorylase,SPAG2

产品验证



Western blot analysis of UAP1 expressed in Mouse brain,Rat brain,HepG2 using UAP1 Rabbit mAb at 1:10 00. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn (<https://www.ablybio.cn/>www.ablybio.cn)