

— ABLYBIO, Help Your Research



MGEA5 Rabbit mAb

货号: **AYM30692**

产品信息

反应	Human,Mouse
宿主	Rabbit
克隆性	Monoclonal
预测反应	
应用	WB IHC IP
推荐浓度	WB: 1:500 - 1:2000 IHC: 1:50 - 1:200 IP: 1:20 - 1:50
理论分子量	76kDa/95kDa/96kDa/102kDa
实测分子量	130kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.75% BSA,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	U-87MG,HeLa
细胞定位	Cytoplasm,Nucleus
纯化	Affinity purification


抗原信息

抗原信息	Recombinant fusion protein corresponding to Human MGEA5.
------	--

靶点信息

研究背景	The dynamic modification of cytoplasmic and nuclear proteins by O-linked N-acetylglucosamine (O-GlcNAc) addition and removal on serine and threonine residues is catalyzed by OGT (MIM 300255), which adds O-GlcNAc, and MGEA5, a glycosidase that removes O-GlcNAc modifications (Gao et al., 2001 [PubMed 11 148210]).[supplied by OMIM, Mar 2008]
基因ID	10724
基因名	MGEA5
Swiss	O60502
别名	MGEA5,MGEA5 Rabbit mAb,Beta-N-acetylglucosaminidase,Beta-N-acetylhexosaminidase,Beta-hexosaminidase,Meningioma-expressed antigen 5,N-acetyl-beta-D-glucosaminidase,N-acetyl-beta-glucosaminidase,Nuclear cytoplasmic O-GlcNAcase and acetyltransferase,HEXC

产品验证



Western blot analysis of MGEA5 expressed in U-87MG,HeLa using MGEA5 Rabbit mAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn (<https://www.ablybio.cn/>www.ablybio.cn)