— ABLYBIO, Help Your Research

p38 MAPK Rabbit mAb

货号: AYM30695

产品信息

反应	Human,Mouse,Rat
宿主	Rabbit
克隆性	Monoclonal
预测反应	
应用	WB IF/ICC
推荐浓度	**WB:** 1:500 - 1:2000 **IF/ICC:** 1:50 - 1:200
理论分子量	29kDa/34kDa/35kDa/41kDa
实测分子量	38kDa
形式	Liquid
保存条件	Store at -20℃. Avoid freeze / thaw cycles. Buffer: PBS with 0.75% BSA,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	HeLa,BxPC-3,Mouse testis,Rat kidney
细胞定位	Cytoplasm,Nucleus
纯化	Affinity purification


抗原信息

抗原信息	Recombinant fusion protein corresponding to Human p38 MAPK.

靶点信息

研究背景	The protein encoded by this gene is a member of the MAP kinase family. MAP kinases act as an integration point for multiple biochemical signals, and are involved in a wide variety of cellular processes such as proliferation, differentiation, transcription regulation and development. This kinase is activated by various environmental stresses and proinflammatory cytokines. The activation requires its phosphorylation by MAP kinase kinases (MKKs), or its autophosphorylation triggered by the interaction of MAP3K7IP1/TAB1 protein with this kinase. The substrates of this kinase include transcription regulator ATF2, MEF2C, and MAX, cell cycle regulator CDC25B, and tumor suppressor p53, which suggest the roles of this kinase in stress related transcription and cell cycle regulation, as well as in genotoxic stress response. Four alternatively spliced transcript variants of this gene encoding distinct isoforms have been reported.
基因ID	1432
基因名	MAPK14
Swiss	Q16539 (https://www.uniprot.org/uniprotkb/Q16539/entry)
别名	p38 MAPK,p38 MAPK Rabbit mAb,MAPK14,Cytokine suppressive anti-inflammatory drug-binding protein,MAP kinase MXI2,MAX-interacting protein 2,Mitogen-activated protein kinase p38 alpha,Stress-activated protein kinase 2a,CSBP,CSBP1,CSBP2,CSPB1,MXI2,SAPK2A

产品验证



Western blot analysis of p38 MAPK expressed in Mouse testis,Rat kidney,HeLa using p38 MAPK Rabbit mAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn (https://www.ablybio.cn//www.ablybio.cn)