— ABLYBIO, Help Your Research

USP22 Rabbit mAb

货号: AYM30704

产品信息

反应	Human,Mouse,Rat
宿主	Rabbit
克隆性	Monoclonal
预测反应	
应用	WB IF/ICC IP
推荐浓度	**WB:** 1:500 - 1:2000 **IF/ICC:** 1:50 - 1:200 **IP:** 1:20 - 1:50
理论分子量	58kDa/59kDa
实测分子量	60kDa
形式	Liquid
保存条件	Store at -20℃. Avoid freeze / thaw cycles. Buffer: PBS with 0.75% BSA,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	U-87MG,293T
细胞定位	Nucleus
纯化	Affinity purification


抗原信息

抗原信息	Recombinant fusion protein corresponding to Human USP22.

靶点信息

研究背景	Histone deubiquitinating component of the transcription regulatory histone acetylation (HAT complex SAGA. Catalyzes the deubiquitination of both histones H2A and H2B, thereby acting as a coactivator. Recruited to specific gene promoters by activators such as MYC, where it is required for transcription. Required for nuclear receptor-mediated transactivation and cell cycle progression.
基因ID	23326
基因名	USP22
Swiss	Q9UPT9 (https://www.uniprot.org/uniprotkb/Q9UPT9/entry)
别名	USP22,USP22 Rabbit mAb,Deubiquitinating enzyme 22,Ubiquitin thioesterase 22,Ubiquitin-specific-processing protease 22,KIAA1063,USP3L

产品验证



Western blot analysis of USP22 expressed in U-87MG,293T using USP22 Rabbit mAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn (https://www.ablybio.cn//www.ablybio.cn)