

DGAT1 Rabbit mAb

货号: **AYM30719**

产品信息

反应	Human
宿主	Rabbit
克隆性	Monoclonal
预测反应	
应用	WB IF/ICC IP FC
推荐浓度	WB: 1:500 - 1:2000 IF/ICC: 1:50 - 1:200 IP: 1:20 - 1:50 FC: 1:20 - 1:50
理论分子量	55kDa
实测分子量	48kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.75% BSA,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	HepG2,A-431,HL-60,Mouse liver
细胞定位	Endoplasmic reticulum membrane,Multi-pass membrane protein
纯化	Affinity purification

抗原信息


抗原信息	Recombinant fusion protein corresponding to Human DGAT1.
序列	AHTILFLKLFSDVNSWCRRARAKAASAGKKASSAAAPHTVSYPDNLTYRDLYYFLFAPTLCYELNFPSPRIRKRFLRRIL EMLFFTQLQVGLIQWWM

靶点信息

研究背景	This gene encodes an multipass transmembrane protein that functions as a key metabolic enzyme. The encoded protein catalyzes the conversion of diacylglycerol and fatty acyl CoA to triacylglycerol. This enzyme can also transfer acyl CoA to retinol. Activity of this protein may be associated with obesity and other metabolic diseases.
------	---

基因ID	8694
基因名	DGAT1
Swiss	O75907
别名	DGAT1;ARAT;ARGP1;DGAT;DIAR7

产品验证



Western blot analysis of DGAT1 expressed in HepG2,A-431,HL-60 using DGAT1 Rabbit mAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn