

HTSF1 Rabbit mAb

货号: **AYM30726**

产品信息

反应	Human,Mouse
宿主	Rabbit
克隆性	Monoclonal
预测反应	
应用	WB IHC IF/ICC IP
推荐浓度	WB: 1:500 - 1:2000 IHC: 1:50 - 1:200 IF/ICC: 1:50 - 1:200 IP: 1:20 - 1:50
理论分子量	85kDa
实测分子量	140kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.75% BSA,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	HeLa,Jurkat,HepG2
细胞定位	Nucleus
纯化	Affinity purification

抗原信息


抗原信息	Recombinant fusion protein corresponding to Human HTSF1.
序列	MSGTNLDGNDEFDEQLRMQELYGDGKDGDTQTDAGGEPDSLGGQPTDTPYEWDLDDKAWFPKITEDFIATYQANYGF SNDGASSSTANVEDVHARTAEPPQEKAPEPTDARKKGEKRKAESGWFWHVEEDRNTNVVYVSGLPPDITVDEFIQLMSKF GIIMRDPQTEEFKVKLYKDNQGNL

靶点信息

研究背景	The protein encoded by this gene functions as a cofactor for the stimulation of transcriptional elongation by HIV-1 Tat, which binds to the HIV-1 promoter through Tat-TAR interaction. This protein may also serve as a dual-function factor to couple transcription and splicing and to facilitate their reciprocal activation. A alternatively spliced transcript variants have been found for this gene.
------	--

基因ID	27336
基因名	HTATSF1
Swiss	O43719
别名	HTATSF1;TAT-SF1;TATSF1;dJ196E23.2

产品验证



Western blot analysis of HTSF1 expressed in HeLa, Jurkat, HepG2 using HTSF1 Rabbit mAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn