

# FDX1 / ADX Rabbit mAb

货号: **AYM30727**

## 产品信息

反应	Human,Mouse,Rat
宿主	Rabbit
克隆性	Monoclonal
预测反应	
应用	WB IHC IF/ICC
推荐浓度	<b>WB:</b> 1:500 - 1:2000 <b>IHC:</b> 1:50 - 1:200 <b>IF/ICC:</b> 1:50 - 1:200
理论分子量	19kDa
实测分子量	13kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.75% BSA,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	HepG2,A-549,U-87MG,Mouse liver,Mouse testis,Mouse kidney,Rat testis,Rat kidney
细胞定位	Mitochondrion matrix
纯化	Affinity purification

## 抗原信息


抗原信息	Recombinant fusion protein corresponding to Human FDX1 / ADX.
序列	MAAAGGARLLRAASAVLGGPAGRWLHHAGSRAGSSGLLRNRGPGGSAEASRSLSVSARARSSSEDKITVHFINRDGETL TTKGKVGDSLDDVVVENNLDI

## 靶点信息

研究背景	This gene encodes a small iron-sulfur protein that transfers electrons from NADPH through ferredoxin reductase to mitochondrial cytochrome P450, involved in steroid, vitamin D, and bile acid metabolism. Pseudogenes of this functional gene are found on chromosomes 20 and 21.
基因ID	2230

基因名	FDX1
Swiss	P10109
别名	FDX1;ADX;FDX;LOH11CR1D

## 产品验证



Western blot analysis of FDX1 / ADX expressed in Mouse liver,Rat testis,HepG2 using FDX1 / ADX Rabbit mAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

## 实验步骤

访问官网浏览详情: [www.ablybio.cn](http://www.ablybio.cn)