

FHL2 Rabbit mAb

货号: **AYM30746**

产品信息

| | |
|-------|---|
| 反应 | Human,Mouse,Rat |
| 宿主 | Rabbit |
| 克隆性 | Monoclonal |
| 预测反应 | |
| 应用 | WB IF/ICC IP |
| 推荐浓度 | WB: 1:500 - 1:2000 IF/ICC: 1:50 - 1:200 IP: 1:20 - 1:50 |
| 理论分子量 | 32kDa |
| 实测分子量 | 32kDa |
| 形式 | Liquid |
| 保存条件 | Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.75% BSA,50% glycerol,pH7.3. |
| 偶联物 | Unconjugated |
| 阳性对照 | Mouse heart,Rat heart,SKOV3,RD |
| 细胞定位 | Cytoplasm,Nucleus |
| 纯化 | Affinity purification |


抗原信息

| | |
|------|--|
| 抗原信息 | Recombinant fusion protein corresponding to Human FHL2. |
| 序列 | DCKDLSYKDRHWHEACFHCSQCRNSLVDKPFPAKEDQLLCTDCYSNEYSSKCQECKKTIMPGTRKMEYKGSSWHETCFI CHRCQQPIGTSFIPKDNQNFVPCYEKQHAMQCVQCKPITTTGGVITYREQPWHKECFVCTACRKQLSGQRFTARDDFA YCLNCFCDLYAKKCAGCTNPISGLGGTKYISFEERQWHNDCFNCKKCSLSLVG |

靶点信息

| | |
|-------|--|
| 研究背景 | This gene encodes a member of the four-and-a-half-LIM-only protein family. Family members contain two highly conserved, tandemly arranged, zinc finger domains with four highly conserved cysteines binding a zinc atom in each zinc finger. This protein is thought to have a role in the assembly of extracellular membranes. Also, this gene is down-regulated during transformation of normal myoblasts to rhabdomyosarcoma cells and the encoded protein may function as a link between presenilin-2 and an intracellular signaling pathway. Multiple alternatively spliced variants encoding different isoforms have been identified. [provided by RefSeq, Jan 2016] |
| 基因ID | 2274 |
| 基因名 | FHL2 |
| Swiss | Q14192 |
| 别名 | AAG11; DRAL; FHL-2; SLIM-3; SLIM3 |

产品验证



Western blot analysis of FHL2 expressed in Mouse heart, Rat heart, SKOV3 using FHL2 Rabbit mAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn