

AHA1 Rabbit mAb

货号: **AYM30783**

产品信息

| | |
|-------|---|
| 反应 | Human,Mouse,Rat |
| 宿主 | Rabbit |
| 克隆性 | Monoclonal |
| 预测反应 | |
| 应用 | WB IP |
| 推荐浓度 | WB: 1:500 - 1:2000 IP: 1:20 - 1:50 |
| 理论分子量 | 43kDa |
| 实测分子量 | 43kDa |
| 形式 | Liquid |
| 保存条件 | Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.75% BSA,50% glycerol,pH7.3. |
| 偶联物 | Unconjugated |
| 阳性对照 | HeLa,293T,RD,Mouse brain,Rat brain |
| 细胞定位 | Cytoplasm,Endoplasmic reticulum,cytosol |
| 纯化 | Affinity purification |

抗原信息


| | |
|------|---|
| 抗原信息 | Recombinant fusion protein corresponding to Human AHA1. |
|------|---|

靶点信息

| | |
|------|--|
| 研究背景 | AHA-1 stimulates the inherent ATPase activity of yeast and human HSP 90 and interacts with the cytoplasmic tail of vesicular stomatitis virus glycoprotein. AHA-1 regulates HSP 90 by influencing the conformational state of the "ATP lid" and consequent N-terminal dimerization. It is crucial for cell viability under non-optimal growth conditions when HSP 90 levels are limiting. AHA-1 is a cytosolic protein and may transiently interact with the endoplasmic reticulum. It can have an affect on one step in the endoplasmic to Golgi trafficking. AHA-1 is expressed in numerous tissues, including brain, heart, skeletal muscle and kidney and , at lower levels, in liver and placenta. It is induced by heat shock and treatment with the HSP 90 inhibitor 17-demeth-oxygeldanamycin. |
| 基因ID | 10598 |

| | |
|-------|--------|
| 基因名 | AHSA1 |
| Swiss | O95433 |
| 别名 | AHA1 |

产品验证



Western blot analysis of AHA1 expressed in Mouse brain,Rat brain,HeLa using AHA1 Rabbit mAb at 1:10 00. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blotting buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn