

Olig3 Rabbit mAb

货号: **AYM30807**

产品信息

| | |
|-------|---|
| 反应 | Human |
| 宿主 | Rabbit |
| 克隆性 | Monoclonal |
| 预测反应 | |
| 应用 | WB IF/ICC |
| 推荐浓度 | WB: 1:500 - 1:2000 IF/ICC: 1:50 - 1:200 |
| 理论分子量 | 29kDa |
| 实测分子量 | 29kDa |
| 形式 | Liquid |
| 保存条件 | Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.75% BSA, 50% glycerol, pH7.3. |
| 偶联物 | Unconjugated |
| 阳性对照 | U-251MG |
| 细胞定位 | Nucleus |
| 纯化 | Affinity purification |

抗原信息


| | |
|------|--|
| 抗原信息 | Recombinant fusion protein corresponding to Human Olig3. |
| 序列 | HAANSVHPVHPILGGALSSGNASSPLSAASLPAIGTIRPPHSLLKAPSTPPALQLGSGFQHWAGLPCPTICMPPPPHLSA LSTANMARLSAESKDLLK |

靶点信息

| | |
|------|--|
| 研究背景 | May determine the distinct specification program of class A neurons in the dorsal part of the spinal cord and suppress specification of class B neurons. |
| 基因ID | 167826 |
| 基因名 | OLIG3 |

| | |
|-------|--|
| Swiss | Q7RTU3 |
| 别名 | OLIG3; Bhlhb7; bHLHe20; oligodendrocyte transcription factor 3 |

产品验证



Western blot analysis of Olig3 expressed in U-251MG using Olig3 Rabbit mAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn