

# MALT1 Rabbit mAb

货号: **AYM30819**

## 产品信息

|       |   |
|-------|---|
| 反应    | Human   |
| 宿主    | Rabbit  |
| 克隆性   | Monoclonal  |
| 预测反应  |   |
| 应用    | WB IF/ICC FC  |
| 推荐浓度  | <b>WB:</b> 1:500 - 1:2000<br><b>IF/ICC:</b> 1:50 - 1:200<br><b>FC:</b> 1:20 - 1:50            |
| 理论分子量 | 91kDa/92kDa   |
| 实测分子量 | 92kDa   |
| 形式    | Liquid  |
| 保存条件  | Store at -20°C. Avoid freeze / thaw cycles.<br>Buffer: PBS with 0.75% BSA,50% glycerol,pH7.3. |
| 偶联物   | Unconjugated  |
| 阳性对照  | HeLa,293T   |
| 细胞定位  | Cytoplasm,Nucleus,perinuclear region  |
| 纯化    | Affinity purification   |


## 抗原信息

|      |   |
|------|---|
| 抗原信息 | Recombinant fusion protein corresponding to Human MALT1 .   |
| 序列   | PGIKITVNPESKAVLAGQFVKLCCRATGHPFVQYQWFKMNKEIPNGNTSELIFNAVHVKDGAFYVCRVNNNFTFEFSQWS<br>QLDVCDIPESFQRSVDGVSESKLQICVEPTSQKLMPGSTLVLCQVAVGSPIPHYQWFKNELPLTHETKKLYMVPYVDLEHQ<br>GTYWCHVYNDRDSQDSKKVEIIDEIENNLGHDPNKEQTTDQPLAKDKVALLIGNMNYREHPKPKAPLVDVYELTNLLRQLD<br>FKVVSLLDLTEYEMRNAVDEFLLLLDKGVYGLLYAGHGYENFGNSFMVPVDAPNPYRSENCLCVQNILKLMQEKETGLNV<br>FLL |

## 靶点信息

|       |   |
|-------|---|
| 研究背景  | This gene has been found to be recurrently rearranged in chromosomal translocation with two other genes - baculoviral IAP repeat-containing protein 3 (also known as apoptosis inhibitor 2) and immunoglobulin heavy chain locus - in mucosa-associated lymphoid tissue lymphomas. The protein encoded by this gene may play a role in NF-kappaB activation. Two alternatively spliced transcript variants encoding different isoforms have been described for this gene. |
| 基因ID  | 10892   |
| 基因名   | MALT1   |
| Swiss | Q9UDY8  |
| 别名    | MALT1;IMD12;MLT;MLT1;PCASP1   |

## 产品验证



Western blot analysis of MALT1 expressed in HeLa,293T using MALT1 Rabbit mAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

## 实验步骤

访问官网浏览详情: [www.ablybio.cn](http://www.ablybio.cn)