

Phospho-JAK2 (Y1007 + Y1008) Rabbit mAb

货号: **AYM30840**

产品信息

反应	Human,Mouse,Rat
宿主	Rabbit
克隆性	Monoclonal
预测反应	
应用	WB IHC ICC IP
推荐浓度	WB: 1:500 - 1:2000 IHC: 1:50 - 1:200 ICC: 1:50 - 1:200 IP: 1:20 - 1:50
理论分子量	130kDa
实测分子量	130kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.75% BSA,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	HT29
细胞定位	Cytoplasm,Endomembrane system,Nucleus,Peripheral membrane protein
纯化	Affinity purification

抗原信息


抗原信息	Recombinant fusion protein corresponding to Human Phospho-JAK2 (Y1007 + Y1008).
序列	KEYYK

靶点信息


研究背景	This gene product is a protein tyrosine kinase involved in a specific subset of cytokine receptor signaling pathways. It has been found to be constitutively associated with the prolactin receptor and is required for responses to gamma interferon. Mice that do not express an active protein for this gene exhibit embryonic lethality associated with the absence of definitive erythropoiesis.
基因ID	3717

基因名	JAK2
Swiss	O60674
别名	JAK2;JTK10;THCYT3

产品验证



Immunohistochemical analysis of paraffin-embedded human cervix cancer, using Phospho-JAK2 (Y1007 + Y1008) Antibody.



Western blot analysis of Phospho-JAK2 (Y1007 + Y1008) expressed in HT29 using Phospho-JAK2 (Y1007 + Y1008) Rabbit mAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn