— ABLYBIO, Help Your Research

Histone H3 (di methyl K9) Rabbit mAb

货号: AYM30898

产品信息

反应	Human,Mouse,Rat
宿主	Rabbit
克隆性	Monoclonal
预测反应	
应用	WB ICC/IF
推荐浓度	**WB:** 1:500 - 1:2000 **IF/ICC:** 1:50 - 1:200
理论分子量	15kDa
实测分子量	15kDa
形式	Liquid
保存条件	Store at -20℃. Avoid freeze / thaw cycles. Buffer: PBS with 0.75% BSA,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	C2C12 cells, HEK-293T cells, THP-1 cells, HeLa cells, HepG2 cells, HSC-T6 cells
细胞定位	Chromosome,Nucleus
纯化	Affinity purification


抗原信息

抗原信息	Recombinant fusion protein corresponding to Human Histone H3 (di methyl K9).


靶点信息

研究背景	Histones are basic nuclear proteins that are responsible for the nucleosome structure of the chromosomal fiber in eukaryotes. This structure consists of approximately 146 bp of DNA wrapped around a nucleosome, an octamer composed of pairs of each of the four core histones (H2A, H2B, H3, and H4). The chromatin fiber is further compacted through the interaction of a linker histone, H1, with the DNA between the nucleosomes to form higher order chromatin structures. This gene is intronless and encodes a replication-dependent histone that is a member of the histone H3 family. Transcripts from this gene lack polyA tails; instead, they contain a palindromic termination element. This gene is found in the large histone gene cluster on chromosome 6p22-p21.3.
基因ID	8350
基因名	H3C1, H3C2, H3C3, H3C4, H3C6, H3C7, H3C8, H3C10, H3C11, H3C12
Swiss	P68431 (https://www.uniprot.org/uniprotkb/P68431/entry)
别名	Histone H3 (di methyl K9),Histone H3 (di methyl K9) Rabbit mAb,H3C1,H3C2,H3C3,H3C4,H3C6,H3C7,H3C8,H3C10,H3C11,H3C12,Histone H3/a,Histone H3/b,Histone H3/c,Histone H3/d,Histone H3/f,Histone H3/h,Histone H3/i,Histone H3/j,Histone H3/k,Histone H3/l,H3FA,HIST1H3A,H3FL,HIST1H3B

产品验证



Western blot analysis of Histone H3 (di methyl K9) expressed in C2C12 cells,HEK-293T cells,THP-1 cells using Histone H3 (di methyl K9) Rabbit mAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.



Immunofluorescence analysis of HeLa cells labelling Histone H3 (di methyl K9) with . The cells were fixed with cold 100% methanol (10min, 4℃) and blocked in 1% BSA/10% normal goat serum/0.3M glycine in 0.1% PBS-Tween 20 for 1h. The cells were then incubate with Goat Anti-Rabbit IgG Fluor488 (1:200，shown in green) at room temprature for 1h.

实验步骤

访问官网浏览详情: www.ablybio.cn (https://www.ablybio.cn//www.ablybio.cn)