

IRAKM Rabbit mAb

货号: **AYM30924**

产品信息

反应	Human
宿主	Rabbit
克隆性	Monoclonal
预测反应	
应用	WB IP
推荐浓度	WB: 1:500 - 1:2000 IP: 1:20 - 1:50
理论分子量	60kDa/67kDa
实测分子量	68kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.75% BSA, 50% glycerol, pH7.3.
偶联物	Unconjugated
阳性对照	THP-1, NCI-H125, Rat liver
细胞定位	cytoplasm, nucleus, plasma membrane
纯化	Affinity purification

抗原信息


抗原信息	Recombinant fusion protein corresponding to Human IRAKM.
序列	VLTGCRVVLDDPKHIQLRDLLRELMEKRGLDSCLSFLDKKVPVPCPRNFSAKLFCLAGRCAATRAKLRPSMDEVLTLESTQASLYFAEDPPTSLKSFRCPSPLFLENVPSIPVEDDESQNNLLPSDEGLRIDRMTQKTPFECSQSEVMFLSLDKKPESKRNEEACNMPSSSCEESWFPKYIVPSQDLRPYKVNIDPSSEAPGHSCRSRPVESSCSSKFSWDEYEQYKKE

靶点信息

研究背景	This gene encodes a member of the interleukin-1 receptor-associated kinase protein family. Members of this family are essential components of the Toll/IL-R immune signal transduction pathways. This protein is primarily expressed in monocytes and macrophages and functions as a negative regulator of Toll-like receptor signaling. Mutations in this gene are associated with a susceptibility to asthma. Alternate splicing results in multiple transcript variants.
------	---

基因ID	11213
基因名	IRAK3
Swiss	Q9Y616
别名	IRAK3;ASRT5;IRAKM

产品验证



Western blot analysis of IRAKM expressed in THP-1, NCI-H125, Rat liver using IRAKM Rabbit mAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn