— ABLYBIO, Help Your Research

CDK1 Rabbit mAb

货号: AYM30936

产品信息

反应	Human
宿主	Rabbit
克隆性	Monoclonal
预测反应	
应用	WB IHC IP ICC/IF
推荐浓度	**WB:** 1:500 - 1:2000 **IHC:** 1:50 - 1:200 **IF/ICC:** 1:50 - 1:200 **IP:** 1:20 - 1:50
理论分子量	27kDa/34kDa
实测分子量	34kDa
形式	Liquid
保存条件	Store at -20℃. Avoid freeze / thaw cycles. Buffer: PBS with 0.75% BSA,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	HT-29,HeLa,NIH/3T3,C6
细胞定位	Cytoplasm,Mitochondrion,Nucleus,centrosome,cytoskeleton,microtubule organizing center,spindle
纯化	Affinity purification


抗原信息

抗原信息	Recombinant fusion protein corresponding to Human CDK1.


靶点信息

研究背景	The protein encoded by this gene is a member of the Ser/Thr protein kinase family. This protein is a catalytic subunit of the highly conserved protein kinase complex known as M-phase promoting factor (MPF), which is essential for G1/S and G2/M phase transitions of eukaryotic cell cycle. Mitotic cyclins stably associate with this protein and function as regulatory subunits. The kinase activity of this protein is controlled by cyclin accumulation and destruction through the cell cycle. The phosphorylation and dephosphorylation of this protein also play important regulatory roles in cell cycle control. Alternatively spliced transcript variants encoding different isoforms have been found for this gene.
基因ID	983
基因名	CDK1
Swiss	P06493 (https://www.uniprot.org/uniprotkb/P06493/entry)
别名	CDK1,CDK1 Rabbit mAb,Cell division control protein 2 homolog,Cell division protein kinase 1,p34 protein kinase,CDC2,CDC28A,CDKN1,P34CDC2


产品验证



Western blot analysis of CDK1 expressed in HT-29,HeLa,NIH/3T3 using CDK1 Rabbit mAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.



Immunohistochemical analysis of paraffin-embedded human colon, using CDK1 Antibody.



Immunofluorescence analysis of ACHN cells labelling CDK1 with . The cells were fixed with cold 100% methanol (10min, 4℃) and blocked in 1% BSA/10% normal goat serum/0.3M glycine in 0.1% PBS-Tween 20 for 1h. The cells were then incubate with Goat Anti-Rabbit IgG Fluor488 (1:200，shown in green) at room temprature for 1h.

实验步骤

访问官网浏览详情: www.ablybio.cn (https://www.ablybio.cn//www.ablybio.cn)