

FKBP38 Rabbit mAb

货号: **AYM30974**

产品信息

反应	Human
宿主	Rabbit
克隆性	Monoclonal
预测反应	
应用	WB IF/ICC FC
推荐浓度	WB: 1:500 - 1:2000 IF/ICC: 1:50 - 1:200 FC: 1:20 - 1:50
理论分子量	27kDa/44kDa
实测分子量	52kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.75% BSA,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	Jurkat,THP-1,BT-474,HeLa,HL-60,Mouse liver,Mouse brain,Mouse eye,Rat brain
细胞定位	Cytoplasmic side,Mitochondrion membrane,Single-pass membrane protein
纯化	Affinity purification

抗原信息


抗原信息	Recombinant fusion protein corresponding to Human FKBP38.
序列	QPPAEEAEQPGALAREFLAAMEPEPAPAPAPEEWLDILGNLLRKKTLVPGPPGSSRPVKGQVVTVHLQTSLENGTRVQEE PELVFTLGDCDVIQALDLSVPLMDVGETAMVTADSKYCYGPQGSRSPIPPHAALCLEVTLKTAVDGPDLEMLTGQERVAL ANRKRECGNAHYQRADFVLAANSYDLAIKAITSSAKVDMTFEEEEALLQ

靶点信息

研究背景	The protein encoded by this gene is a member of the immunophilin protein family, which play a role in immunoregulation and basic cellular processes involving protein folding and trafficking. Unlike the other members of the family, this encoded protein does not seem to have PPIase/rotamase activity. It may have a role in neurons associated with memory function.
------	--

基因ID	23770
基因名	FKBP8
Swiss	Q14318
别名	FKBP8;FKBP38;FKBPr38

产品验证



Western blot analysis of FKBP38 expressed in Jurkat, THP-1, BT-474 using FKBP38 Rabbit mAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn