

NRCAM Rabbit mAb

货号: **AYM30977**

产品信息

反应	Human
宿主	Rabbit
克隆性	Monoclonal
预测反应	
应用	WB IP
推荐浓度	WB: 1:500 - 1:2000 IP: 1:20 - 1:50
理论分子量	144kDa
实测分子量	144kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.75% BSA,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	SH-SY5Y,Mouse spinal cord,Mouse brain
细胞定位	axon,external side of plasma membrane,extracellular region,glutamatergic synapse,neuron projection,plasma membrane,postsynaptic density membrane
纯化	Affinity purification

抗原信息


抗原信息	Recombinant fusion protein corresponding to Human NRCAM.
------	--

靶点信息

研究背景	Cell adhesion molecules (CAMs) are members of the immunoglobulin superfamily. This gene encodes a neuronal cell adhesion molecule with multiple immunoglobulin-like C2-type domains and fibronectin type-III domains. This ankyrin-binding protein is involved in neuron-neuron adhesion and promotes directional signaling during axonal cone growth. This gene is also expressed in non-neural tissues and may play a general role in cell-cell communication via signaling from its intracellular domain to the actin cytoskeleton during directional cell migration. Allelic variants of this gene have been associated with autism and addiction vulnerability. Alternative splicing results in multiple transcript variants encoding different isoforms.
基因ID	4897

基因名	NRCAM
Swiss	Q92823
别名	NRCAM

产品验证



Western blot analysis of NRCAM expressed in SH-SY5Y, Mouse spinal cord, Mouse brain using NRCAM Rabbit mAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30 ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn