

RAGE Rabbit mAb

货号: **AYM30978**

产品信息

反应	Human,Mouse,Rat
宿主	Rabbit
克隆性	Monoclonal
预测反应	
应用	WB
推荐浓度	WB: 1:500 - 1:2000
理论分子量	13kDa/35-44kDa
实测分子量	48kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.75% BSA,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	A-549,NCI-H460,Mouse small intestine,Rat kidney
细胞定位	Cell membrane,Secreted,Single-pass type I membrane protein
纯化	Affinity purification

抗原信息


抗原信息	Recombinant fusion protein corresponding to Human RAGE.
序列	IGEPLVLKCKGAPKKPPQRLEWKLNTGRTEAWKVLSPQGGGPWDSVARVLPNGSLFLPAVGIQDEGIFRCQAMNRNGKE TKSNYRVRVYQIPGKPEIVDSASELTAGVPNKVGTVCVSEGSYPAGTLSWHLDGKPLVPNEKGVSVKEQTRRHHPETGLFTLQ SELMVTPARGGDPRTFSCSFSPGLPRHRALRTAPIQPRVWEPVPLEEVQLVVEPEGGAVAPGGTVTLTCEVPAQPSPIH WMKDGVPLPLPPSPVLILPEIGPQDQGTYSVATHSSHGPGQESRAVSISIIIEPGEEGPTA

靶点信息

研究背景	The advanced glycosylation end product (AGE) receptor encoded by this gene is a member of the immunoglobulin superfamily of cell surface receptors. It is a multiligand receptor, and besides AGE, interacts with other molecules implicated in homeostasis, development, and inflammation, and certain diseases, such as diabetes and Alzheimer's disease. Many alternatively spliced transcript variants encoding different isoforms, as well as non-protein-coding variants, have been described for this gene (PMID:18089847).
------	--

基因ID	177
基因名	AGER
Swiss	Q15109
别名	AGER;RAGE;SCARJ1

产品验证



Western blot analysis of RAGE expressed in Mouse small intestine,Rat kidney,A-549 using RAGE Rabbit m Ab at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn