

# Cytokeratin 17 Rabbit mAb

货号: **AYM30984**

## 产品信息

反应	Human,Mouse,Rat
宿主	Rabbit
克隆性	Monoclonal
预测反应	
应用	WB IHC IF/ICC FC
推荐浓度	<b>WB:</b> 1:500 - 1:2000 <b>IHC:</b> 1:50 - 1:200 <b>IF/ICC:</b> 1:50 - 1:200 <b>FC:</b> 1:20 - 1:50
理论分子量	48kDa
实测分子量	48kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.75% BSA,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	A-431,HeLa,A-549 negative,Mouse stomach
细胞定位	Cytoplasm
纯化	Affinity purification

## 抗原信息


抗原信息	Recombinant fusion protein corresponding to Human Cytokeratin 17.
序列	NRVCVQLSQQGLIGSVEEQLAQLRCMEQQNQEQYKILLDVKTRLEQEIATYRRLEGEDAHLTQYKKEPVTTTRQVRTIVEEV QDGKVISSREQVHQTTTR

## 靶点信息


研究背景	This gene encodes the type I intermediate filament chain keratin 17, expressed in nail bed, hair follicle, sebaceous glands, and other epidermal appendages. Mutations in this gene lead to Jackson-Lawler type pachyonychia congenita and steatocystoma multiplex.
基因ID	3872

基因名	KRT17
Swiss	Q04695
别名	KRT17;39.1;CK-17;K17;PC;PC2;PCHC1

## 产品验证



Immunohistochemical analysis of paraffin-embedded human tonsil, using Cytokeratin 17 Antibody.



Western blot analysis of Cytokeratin 17 expressed in Mouse stomach, A-431, HeLa using Cytokeratin 17 Rabbit mAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

## 实验步骤

访问官网浏览详情: [www.ablybio.cn](http://www.ablybio.cn)