

# CD99 Rabbit mAb

货号: **AYM30999**

## 产品信息

反应	Human
宿主	Rabbit
克隆性	Monoclonal
预测反应	
应用	WB IHC ICC IP FC
推荐浓度	<b>WB:</b> 1:500 - 1:2000 <b>IHC:</b> 1:50 - 1:200 <b>ICC:</b> 1:50 - 1:200 <b>IP:</b> 1:20 - 1:50 <b>FC:</b> 1:20 - 1:50
理论分子量	16kDa/17kDa/18kDa
实测分子量	28kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.75% BSA,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	Jurkat,U-251MG,U-937,BT-474,THP-1
细胞定位	Membrane,Single-pass type I membrane protein
纯化	Affinity purification


## 抗原信息

抗原信息	Recombinant fusion protein corresponding to Human CD99.
序列	DGGFDLSDALPDNENKKPTAIPKKPSAGDDFDLGDVAVDGENDDRPPNPPKMPNPNPNHPSSSGFSFDADLADGVS GGEGKGGSDGGGSHRKEGEEADAPGVIPGIVGAVVVAVAGAISSFIAYQKKKLCFKENAEQGEVDMESHNRNANAEPVQ RTLLEK

## 靶点信息

研究背景	The protein encoded by this gene is a cell surface glycoprotein involved in leukocyte migration, T-cell adhesion, ganglioside GM1 and transmembrane protein transport, and T-cell death by a caspase-independent pathway. In addition, the encoded protein may have the ability to rearrange the actin cytoskeleton and may also act as an oncosuppressor in osteosarcoma. This gene is found in the pseudoautosomal region of chromosomes X and Y and escapes X-chromosome inactivation. There is a related pseudogene located immediately adjacent to this locus.
基因ID	4267
基因名	CD99
Swiss	P14209
别名	CD99;HBA71;MIC2;MIC2X;MIC2Y;MSK5X

## 产品验证



Western blot analysis of CD99 expressed in Jurkat,U-251MG,U-937 using CD99 Rabbit mAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

## 实验步骤

访问官网浏览详情: [www.ablybio.cn](http://www.ablybio.cn)