

TNPO3 Rabbit mAb

货号: **AYM31022**

产品信息

反应	Human,Mouse,Rat
宿主	Rabbit
克隆性	Monoclonal
预测反应	
应用	WB IHC IF/ICC
推荐浓度	WB: 1:500 - 1:2000 IHC: 1:50 - 1:200 IF/ICC: 1:50 - 1:200
理论分子量	100kDa
实测分子量	100kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.75% BSA,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	HeLa,RD,Jurkat,Mouse lung,Mouse testis,Mouse thymus,Rat testis
细胞定位	cytoplasm,nuclear envelope
纯化	Affinity purification

抗原信息


抗原信息	Recombinant fusion protein corresponding to Human TNPO3.
序列	QVMNQLGQQLVSQLLHTCCFCLPPYTLPDVAEVLWEIMQVDRPTFCRWLENSLKGLPKETTGVAVTVTHKQLTDFHKQV TSAEECKQVCWALRDFTRLFR

靶点信息

研究背景	The protein encoded by this gene is a nuclear import receptor for serine/arginine-rich (SR) proteins such as the splicing factors SFRS1 and SFRS2. The encoded protein has also been shown to be involved in HIV-1 infection, apparently through interaction with the HIV-1 capsid protein. Two transcript variants encoding different isoforms as well as a noncoding transcript have been found for this gene.[provided by RefSeq, Jul 2010]
------	--

基因ID	23534
基因名	TNPO3
Swiss	Q9Y5L0
别名	IPO12; LGMD1F; MTR10A; TRN-SR; TRN-SR2; TRNSR

产品验证



Western blot analysis of TNPO3 expressed in Mouse lung, Rat testis, HeLa using TNPO3 Rabbit mAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn