

# UBE4A Rabbit mAb

货号: **AYM31026**

## 产品信息

反应	Human
宿主	Rabbit
克隆性	Monoclonal
预测反应	
应用	WB IF/ICC FC
推荐浓度	<b>WB:</b> 1:500 - 1:2000 <b>IF/ICC:</b> 1:50 - 1:200 <b>FC:</b> 1:20 - 1:50
理论分子量	123kDa
实测分子量	123kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.75% BSA,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	HUVEC,Mouse brain,Mouse kidney,Rat liver,Rat brain
细胞定位	Cytoplasm
纯化	Affinity purification

## 抗原信息


抗原信息	Recombinant fusion protein corresponding to Human UBE4A.
序列	MTDQENNNNISSNPFAALFGSLADAKQFAAIQKEQLKQSQDELPAASPDDSDNSVSESLDEFDYSVAEISRSFRSQQEICE QLNINHMIQRIFLITLDNSD

## 靶点信息

研究背景	This gene encodes a member of the U-box ubiquitin ligase family. The encoded protein is involved in mult ubiquitin chain assembly and plays a critical role in chromosome condensation and separation through the polyubiquitination of securin. Autoantibodies against the encoded protein may be markers for scleroderma and Crohn's disease. A pseudogene of this gene is located on the long arm of chromosome 3. Alternatively spliced transcript variants encoding multiple isoforms have been observed for this gene.
------	--

基因ID	9354
基因名	UBE4A
Swiss	Q14139
别名	UBE4A; E4; UBOX2; UFD2; ubiquitination factor E4A

## 产品验证



Western blot analysis of UBE4A expressed in HUVEC, Mouse brain, Mouse kidney using UBE4A Rabbit mAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

## 实验步骤

访问官网浏览详情: [www.ablybio.cn](http://www.ablybio.cn)