

— ABLYBIO, Help Your Research



TRAPPC2 Rabbit mAb

货号: **AYM31031**

产品信息

反应	Human,Mouse,Rat
宿主	Rabbit
克隆性	Monoclonal
预测反应	
应用	WB IHC ICC
推荐浓度	WB: 1:500 - 1:2000 IHC: 1:50 - 1:200 ICC: 1:50 - 1:200
理论分子量	14kDa/16kDa/20kDa
实测分子量	16kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.75% BSA,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	SKOV3,PC-3,SH-SY5Y,Mouse brain,Mouse thymus,Rat brain
细胞定位	Cytoplasm,Endoplasmic reticulum,Golgi apparatus,Nucleus,perinuclear region
纯化	Affinity purification


抗原信息

抗原信息	Recombinant fusion protein corresponding to Human TRAPPC2.
------	--

靶点信息

研究背景	The protein encoded by this gene is thought to be part of a large multi-subunit complex involved in the targeting and fusion of endoplasmic reticulum-to-Golgi transport vesicles with their acceptor compartment. In addition, the encoded protein can bind c-myc promoter-binding protein 1 and block its transcriptional repression capability. Mutations in this gene are a cause of spondyloepiphyseal dysplasia tarda (SED). A processed pseudogene of this gene is located on chromosome 19, and other pseudogenes are found on chromosomes 8 and Y. Alternatively spliced transcript variants have been found for this gene.
基因ID	6399
基因名	TRAPPC2
Swiss	P0D181 (https://www.uniprot.org/uniprotkb/P0D181/entry)
别名	TRAPPC2,TRAPPC2 Rabbit mAb,Sedlin,SEDL

产品验证



Western blot analysis of TRAPPC2 expressed in Mouse brain,Rat brain,SKOV3 using TRAPPC2 Rabbit mAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn (<https://www.ablybio.cn/www.ablybio.cn>)