

# TRAPPC2 Rabbit mAb

货号: **AYM31031**

## 产品信息

反应	Human,Mouse,Rat
宿主	Rabbit
克隆性	Monoclonal
预测反应	
应用	WB IHC ICC
推荐浓度	<b>WB:</b> 1:500 - 1:2000 <b>IHC:</b> 1:50 - 1:200 <b>ICC:</b> 1:50 - 1:200
理论分子量	14kDa/16kDa/20kDa
实测分子量	16kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.75% BSA,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	SKOV3,PC-3,SH-SY5Y,Mouse brain,Mouse thymus,Rat brain
细胞定位	Cytoplasm,Endoplasmic reticulum,Golgi apparatus,Nucleus,perinuclear region
纯化	Affinity purification

## 抗原信息


抗原信息	Recombinant fusion protein corresponding to Human TRAPPC2.
------	--

## 靶点信息

研究背景	The protein encoded by this gene is thought to be part of a large multi-subunit complex involved in the targeting and fusion of endoplasmic reticulum-to-Golgi transport vesicles with their acceptor compartment. In addition, the encoded protein can bind c-myc promoter-binding protein 1 and block its transcriptional repression capability. Mutations in this gene are a cause of spondyloepiphyseal dysplasia tarda (SED). A processed pseudogene of this gene is located on chromosome 19, and other pseudogenes are found on chromosomes 8 and Y. Alternatively spliced transcript variants have been found for this gene.
基因ID	6399

基因名	TRAPPC2
Swiss	PODI81
别名	TRAPPC2

## 产品验证



Western blot analysis of TRAPPC2 expressed in Mouse brain,Rat brain,SKOV3 using TRAPPC2 Rabbit m Ab at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 12 0s.

## 实验步骤

访问官网浏览详情: [www.ablybio.cn](http://www.ablybio.cn)