

NAIP Rabbit mAb

货号: **AYM31040**

产品信息

反应	Human
宿主	Rabbit
克隆性	Monoclonal
预测反应	
应用	WB
推荐浓度	WB: 1:500 - 1:2000
理论分子量	
实测分子量	160kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.75% BSA,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	Rat spleen,Rat lung
细胞定位	basolateral plasma membrane,cytoplasm,inflammasome complex,IPAF inflammasome complex
纯化	Affinity purification


抗原信息

抗原信息	Recombinant fusion protein corresponding to Human NAIP.
序列	LLRSIHFPPIRGNKTSAPRAHFSVLETCTFDKSVPTIDQDYASAFEPMNEWERNLAEKEDNVKSYMMDMQRASPDLSTGYWK LSPKQYKIPCLEVDVNDIDVV

靶点信息

研究背景	This gene is part of a 500 kb inverted duplication on chromosome 5q13. This duplicated region contains at least four genes and repetitive elements which make it prone to rearrangements and deletions. The repetitiveness and complexity of the sequence have also caused difficulty in determining the organization of this genomic region. This copy of the gene is full length; additional copies with truncations and internal deletions are also present in this region of chromosome 5q13. It is thought that this gene is a modifier of spinal muscular atrophy caused by mutations in a neighboring gene, SMN1. The protein encoded by this gene contains regions of homology to two baculovirus inhibitor of apoptosis proteins, and it is able to suppress apoptosis induced by various signals. Alternative splicing and the use of alternative promoters result in multiple transcript variants.
基因ID	4671
基因名	NAIP
Swiss	Q13075
别名	NAIP;BIRC1;NLRB1;psiNAIP

产品验证



Western blot analysis of NAIP expressed in Rat spleen, Rat lung using NAIP Rabbit mAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn