

ARF6 Rabbit mAb

货号: **AYM31048**

产品信息

反应	Human,Mouse,Rat
宿主	Rabbit
克隆性	Monoclonal
预测反应	
应用	WB IF/ICC
推荐浓度	WB: 1:500 - 1:2000 IF/ICC: 1:50 - 1:200
理论分子量	20kDa
实测分子量	20kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.75% BSA,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	293T,HepG2,Mouse brain,Mouse kidney,Rat brain
细胞定位	Cell membrane,Cell projection,Cleavage furrow,Cytoplasm,Endosome membrane,Golgi apparatus,Lipid-anchored,Midbody,Recycling endosome membrane,filopodium membrane
纯化	Affinity purification

抗原信息


抗原信息	Recombinant fusion protein corresponding to Human ARF6.
序列	HYTGTQGLIFVVDCAHDRIDEARQELHRIINDREMRDAILIFANKQDLDPAMKPHEIQEKLGLTRIRDNRWYVQPSCATSGDGLYEGLTWLTSNYKS

靶点信息

研究背景	This gene encodes a member of the human ARF gene family, which is part of the RAS superfamily. The ARF genes encode small guanine nucleotide-binding proteins that stimulate the ADP-ribosyltransferase activity of cholera toxin and play a role in vesicular trafficking and as activators of phospholipase D. The product of this gene is localized to the plasma membrane, and regulates vesicular trafficking, remodeling of membrane lipids, and signaling pathways that lead to actin remodeling. A pseudogene of this gene is located on chromosome 7. [provided by RefSeq, Jul 2008]
------	---

基因ID	382
基因名	ARF6
Swiss	P62330
别名	ARF6; ADP-ribosylation factor 6

产品验证



Western blot analysis of ARF6 expressed in Mouse brain, Rat brain, 293T using ARF6 Rabbit mAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn