

# Flotillin 1 Rabbit mAb

货号: **AYM31060**

## 产品信息

反应	Human,Mouse
宿主	Rabbit
克隆性	Monoclonal
预测反应	
应用	WB
推荐浓度	<b>WB:</b> 1:500 - 1:2000
理论分子量	42kDa/47kDa
实测分子量	48kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.75% BSA,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	293T,HeLa,A-549,Mouse brain,Rat brain
细胞定位	Cell membrane,Endosome,Melanosome,Membrane,Peripheral membrane protein,caveola
纯化	Affinity purification

## 抗原信息


抗原信息	Recombinant fusion protein corresponding to Human Flotillin 1 .
序列	HDDQDYHLHSLGKARTAQVQKDARIGEAERDAGIREAKAKQEKVSAQYLSEIEMAKAQRDYELKKAAYDIEVNTRRAQADLAYQLQVAKTKQQIEEQRVQVQVVERAQQVAVQEIQEIARREKELEARVRKPAAERYKLERLAEAEKSQLIMQAEAEAAASVRMRGEAEFAIGARARAEAEQMAKKAFAFQLYQEAQLDMLLEKLPQVAEEISGPLTSANKITLVSSGSGTMGAAKVTGEVLDILTRLPEVERLTGVSISQVNHKPLRTA

## 靶点信息

研究背景	This gene encodes an protein that localizes to the caveolae, which are small domains on the inner cell membranes. This protein plays a role in vesicle trafficking and cell morphology. Alternative splicing results in multiple transcript variants.
基因ID	10211

基因名	FLOT1
Swiss	O75955
别名	FLOT1

## 产品验证



Western blot analysis of Flotillin 1 expressed in Mouse brain,293T,HeLa using Flotillin 1 Rabbit mAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

## 实验步骤

访问官网浏览详情: [www.ablybio.cn](http://www.ablybio.cn)