

DDB2 Rabbit mAb

货号: **AYM31067**

产品信息

反应	Human
宿主	Rabbit
克隆性	Monoclonal
预测反应	
应用	WB IHC IF/ICC FC
推荐浓度	WB: 1:500 - 1:2000 IHC: 1:50 - 1:200 IF/ICC: 1:50 - 1:200 FC: 1:20 - 1:50
理论分子量	17kDa/26kDa/40kDa/47kDa
实测分子量	45kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.75% BSA,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	Mouse liver,Rat liver
细胞定位	Nucleus
纯化	Affinity purification


抗原信息

抗原信息	Recombinant fusion protein corresponding to Human DDB2.
序列	MAPKKRPETQKTSEIVLRPRNKRSRSPLELEPEAKKLCAGSGPSRRCDSDCLWVGLAGPQILPPCRSIVRTLHQHKLGRAS WPSVQQGLQQSFLHTLDSYRILQKAAPFDRRATSLAWHPTHSTVAVGSKGGDIMLWNFGIKDKPTFIKGIGAGGSITGL KFNPLNTNQFYASSMEGTTRLQDFKGNILRVFASDITINIWFCSLDVSASSRMVVTGDNVGNVILLNMDGKELWNLRMH KKKVTHVALNPCCDWFLATASVDQTVKIWDLRQVRGKASFLYSLPHRHPVNAACFSPDGARLLTTDQKSEIRVY

靶点信息

研究背景	This gene encodes a protein that is necessary for the repair of ultraviolet light-damaged DNA. This protein is the smaller subunit of a heterodimeric protein complex that participates in nucleotide excision repair, and this complex mediates the ubiquitylation of histones H3 and H4, which facilitates the cellular response to DNA damage. This subunit appears to be required for DNA binding. Mutations in this gene cause xeroderma pigmentosum complementation group E, a recessive disease that is characterized by an increased sensitivity to UV light and a high predisposition for skin cancer development, in some cases accompanied by neurological abnormalities. Two transcript variants encoding different isoforms have been found for this gene.
基因ID	1643
基因名	DDB2
Swiss	Q92466
别名	DDB2;DDBB;UV-DDB2;XPE

产品验证



Western blot analysis of DDB2 expressed in Mouse liver,Rat liver using DDB2 Rabbit mAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn