

# HSPA14 Rabbit mAb

货号: **AYM31071**

## 产品信息

反应	Human,Mouse,Rat
宿主	Rabbit
克隆性	Monoclonal
预测反应	
应用	WB IHC IF/ICC IP
推荐浓度	<b>WB:</b> 1:500 - 1:2000 <b>IHC:</b> 1:50 - 1:200 <b>IF/ICC:</b> 1:50 - 1:200 <b>IP:</b> 1:20 - 1:50
理论分子量	55kDa
实测分子量	55kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.75% BSA,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	293T,HepG2,Mouse thymus
细胞定位	Cytoplasm,cytosol
纯化	Affinity purification

## 抗原信息


抗原信息	Recombinant fusion protein corresponding to Human HSPA14.
序列	MAAIGVHLGCTSACVAVYKDGGRAGVVANDAGDRVTPAVVAYSENEEIVGLAAKQSRIRNISNTVMKVQILGRSSSDPQA QKYIAESKCLVIEKNGKRLRY

## 靶点信息

研究背景	Component of the ribosome-associated complex (RAC, a complex involved in folding or maintaining nascent polypeptides in a folding-competent state. In the RAC complex, binds to the nascent polypeptide chain, while DNAJC2 stimulates its ATPase activity.
基因ID	51182

基因名	HSPA14
Swiss	Q0VDF9
别名	HSP70-4; HSP70L1

## 产品验证



Western blot analysis of HSPA14 expressed in 293T, HepG2, Mouse thymus using HSPA14 Rabbit mAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

## 实验步骤

访问官网浏览详情: [www.ablybio.cn](http://www.ablybio.cn)