

IRAK4 Rabbit mAb

货号: **AYM31109**

产品信息

反应	Human
宿主	Rabbit
克隆性	Monoclonal
预测反应	
应用	WB IHC IF/ICC FC
推荐浓度	WB: 1:500 - 1:2000 IHC: 1:50 - 1:200 IF/ICC: 1:50 - 1:200 FC: 1:20 - 1:50
理论分子量	37kDa/51kDa
实测分子量	52kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.75% BSA,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	293T
细胞定位	Cytoplasm
纯化	Affinity purification

抗原信息


抗原信息	Recombinant fusion protein corresponding to Human IRAK4.
序列	MNKPITPSTYVRCLNVGLIRKLSDFIDPQEGWKKLAVAIAIKKPSGDDRYNQFHIRRFEALLQTGKSPTSELLFDWGTTNCTV GDLVDLLIQNEFFAPASLLLLPDAVPKTANTLPSKEAITVQQKQMPFCDKDRITLTPVQNLEQSYMPPDSSSPENKSLEVSD TRFHSFSFYELKNVTNNFDERPISVGGNKMGEVGGV

靶点信息

研究背景	This gene encodes a kinase that activates NF-kappaB in both the Toll-like receptor (TLR) and T-cell receptor (TCR) signaling pathways. The protein is essential for most innate immune responses. Mutations in this gene result in IRAK4 deficiency and recurrent invasive pneumococcal disease. Multiple transcript variants encoding different isoforms have been found for this gene.
------	--

基因ID	51135
基因名	IRAK4
Swiss	Q9NWZ3
别名	IRAK4;IPD1;IRAK-4;NY-REN-64;REN64

产品验证



Western blot analysis of IRAK4 expressed in 293T using IRAK4 Rabbit mAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn