

— ABLYBIO, Help Your Research



# NUP62 Rabbit mAb

货号: **AYM31120**

## 产品信息

反应	Human
宿主	Rabbit
克隆性	Monoclonal
预测反应	
应用	WB
推荐浓度	<b>WB:</b> 1:500 - 1:2000
理论分子量	53kDa
实测分子量	70kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.75% BSA,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	A-549,HeLa,HT-29,Raji,BT-474,U-937
细胞定位	Cytoplasm,Nucleus,cytoskeleton,nuclear pore complex,spindle pole
纯化	Affinity purification


## 抗原信息

抗原信息	Recombinant fusion protein corresponding to Human NUP62.
------	--

## 靶点信息

研究背景	The nuclear pore complex is a massive structure that extends across the nuclear envelope, forming a gateway that regulates the flow of macromolecules between the nucleus and the cytoplasm. Nucleoporins are the main components of the nuclear pore complex in eukaryotic cells. The protein encoded by this gene is a member of the FG-repeat containing nucleoporins and is localized to the nuclear pore central plug. This protein associates with the importin alpha/beta complex which is involved in the import of proteins containing nuclear localization signals. Multiple transcript variants of this gene encode a single protein isoform.
基因ID	23636
基因名	NUP62
Swiss	P37198
别名	NUP62,NUP62 Rabbit mAb,62 kDa nucleoporin,Nucleoporin Nup62

## 产品验证



Western blot analysis of NUP62 expressed in A-549, HeLa, HT-29 using NUP62 Rabbit mAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

## 实验步骤

访问官网浏览详情: [www.ablybio.cn](http://www.ablybio.cn) (<https://www.ablybio.cn/>[www.ablybio.cn](http://www.ablybio.cn))