

MRC2 Rabbit mAb

货号: **AYM31143**

产品信息

反应	Human
宿主	Rabbit
克隆性	Monoclonal
预测反应	
应用	WB
推荐浓度	WB: 1:500 - 1:2000
理论分子量	166kDa
实测分子量	190kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.75% BSA,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	SH-SY5Y,HeLa,Mouse kidney
细胞定位	Membrane,Single-pass type I membrane protein
纯化	Affinity purification

抗原信息


抗原信息	Recombinant fusion protein corresponding to Human MRC2.
序列	RWHCRTLGDQLSLLLGARTSNISKPGTLERGDQTRSGQWRIYGSEEDLCALPYHEVYTIQGNSHGKPCTIPFKYDNQWFH GCTSTGREDGHLWCATTQDYGKDERWGFCEPKSNDCEFWDKDQLTDSCYQFNQSTLSWREAWASCEQQGADLLSI TEIHEQTYINGLLTGYSSTLWIGL

靶点信息

研究背景	This gene encodes a member of the mannose receptor family of proteins that contain a fibronectin type II domain and multiple C-type lectin-like domains. The encoded protein plays a role in extracellular matrix remodeling by mediating the internalization and lysosomal degradation of collagen ligands. Expression of this gene may play a role in the tumorigenesis and metastasis of several malignancies including breast cancer, gliomas and metastatic bone disease.
基因ID	9902

基因名	MRC2
Swiss	Q9UBG0
别名	MRC2;CD280;CLEC13E;ENDO180;UPARAP

产品验证



Western blot analysis of MRC2 expressed in SH-SY5Y, HeLa, Mouse kidney using MRC2 Rabbit mAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn