

# PDLIM1 Rabbit mAb

货号: **AYM31156**

## 产品信息

反应	Human,Mouse,Rat
宿主	Rabbit
克隆性	Monoclonal
预测反应	
应用	WB IF/ICC IP FC
推荐浓度	<b>WB:</b> 1:500 - 1:2000 <b>IF/ICC:</b> 1:50 - 1:200 <b>IP:</b> 1:20 - 1:50 <b>FC:</b> 1:20 - 1:50
理论分子量	36kDa
实测分子量	36kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.75% BSA,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	BxPC-3,THP-1,Mouse thymus
细胞定位	Cytoplasm,cytoskeleton
纯化	Affinity purification

## 抗原信息


抗原信息	Recombinant fusion protein corresponding to Human PDLIM1.
序列	LQEKQELNEPPKQSTSFLVLQEILESEEKGDPNKPSGFRSVKAPVTKVAASIGNAQKLPMDCKCGTGIVGVFVKLRDRHRH PECYVCTDCGTNLKQKGHFF

## 靶点信息

研究背景	This gene encodes a member of the enigma protein family. The protein contains two protein interacting domains, a PDZ domain at the amino terminal end and one to three LIM domains at the carboxyl terminal. It is a cytoplasmic protein associated with the cytoskeleton. The protein may function as an adapter to bring other LIM-interacting proteins to the cytoskeleton. Pseudogenes associated with this gene are located on chromosomes 3, 14 and 17.
------	---

基因ID	9124
基因名	PDLIM1
Swiss	O00151
别名	PDLIM1; CLIM1; CLP-36; CLP36; HEL-S-112; hCLIM1; PDZ and LIM domain 1

## 产品验证



Western blot analysis of PDLIM1 expressed in BxPC-3, THP-1, Mouse thymus using PDLIM1 Rabbit mAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

## 实验步骤

访问官网浏览详情: [www.ablybio.cn](http://www.ablybio.cn)