

— ABLYBIO, Help Your Research



# TRIB3 Rabbit mAb

货号: **AYM31165**

## 产品信息

反应	Human
宿主	Rabbit
克隆性	Monoclonal
预测反应	
应用	WB IP
推荐浓度	<b>WB:</b> 1:500 - 1:2000 <b>IP:</b> 1:20 - 1:50
理论分子量	40kDa
实测分子量	40kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.75% BSA,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	K-562
细胞定位	cytosol,nucleoplasm,nucleus,plasma membrane
纯化	Affinity purification


## 抗原信息

抗原信息	Recombinant fusion protein corresponding to Human TRIB3.
------	--

## 靶点信息

研究背景	The protein encoded by this gene is a putative protein kinase that is induced by the transcription factor NF- $\kappa$ B. The encoded protein is a negative regulator of NF- $\kappa$ B and can also sensitize cells to TNF- $\alpha$ and TRAIL-induced apoptosis. In addition, this protein can negatively regulate the cell survival serine-threonine kinase AKT1. Differential promoter usage and alternate splicing result in multiple transcript variants. [provided by RefSeq, Jul 2014]
基因ID	57761
基因名	TRIB3
Swiss	Q96RU7
别名	TRIB3,TRIB3 Rabbit mAb,Neuronal cell death-inducible putative kinase,SINK,p65-interacting inhibitor of NF- $\kappa$ B,C20orf97,NIPK,SKIP3,TRB3

## 产品验证



Western blot analysis of TRIB3 expressed in K-562 using TRIB3 Rabbit mAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

## 实验步骤

访问官网浏览详情: [www.ablybio.cn](http://www.ablybio.cn) (<https://www.ablybio.cn/www.ablybio.cn>)