— ABLYBIO, Help Your Research

TACC3 Rabbit mAb

货号: AYM31168

产品信息

反应	Human,Mouse,Rat
宿主	Rabbit
克隆性	Monoclonal
预测反应	
应用	WB IHC ICC/IF
推荐浓度	**WB:** 1:500 - 1:2000 **IHC:** 1:50 - 1:200 **IF/ICC:** 1:50 - 1:200
理论分子量	140kDa
实测分子量	140kDa
形式	Liquid
保存条件	Store at -20℃. Avoid freeze / thaw cycles. Buffer: PBS with 0.75% BSA,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	293T
细胞定位	cytoplasm,cytosol,mitotic spindle,spindle pole
纯化	Affinity purification


抗原信息

抗原信息	Recombinant fusion protein corresponding to Human TACC3.


靶点信息

研究背景	This gene encodes a member of the transforming acidic colied-coil protein family. The encoded protein is a motor spindle protein that may play a role in stabilization of the mitotic spindle. This protein may also p lay a role in growth a differentiation of certain cancer cells. [provided by RefSeq, Nov 2011]
基因ID	10460
基因名	TACC3
Swiss	Q9Y6A5 (https://www.uniprot.org/uniprotkb/Q9Y6A5/entry)
别名	TACC3,TACC3 Rabbit mAb,ERIC-1,ERIC1


产品验证



Western blot analysis of TACC3 expressed in 293T using TACC3 Rabbit mAb at 1:1000. Secondary antibod y: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.



Immunohistochemical analysis of paraffin-embedded human liver cancer, using TACC3 Antibody.



Immunofluorescence analysis of HeLa cells labelling TACC3 with . The cells were fixed with cold 100% met hanol (10min, 4℃) and blocked in 1% BSA/10% normal goat serum/0.3M glycine in 0.1% PBS-Tween 20 fo r 1h. The cells were then incubate with Goat Anti-Rabbit IgG Fluor488 (1:200，shown in green) at room te mprature for 1h

实验步骤

访问官网浏览详情: www.ablybio.cn (https://www.ablybio.cn//www.ablybio.cn)