

# TACC3 Rabbit mAb

货号: **AYM31168**

## 产品信息

反应	Human,Mouse,Rat
宿主	Rabbit
克隆性	Monoclonal
预测反应	
应用	WB IHC IF/ICC
推荐浓度	<b>WB:</b> 1:500 - 1:2000 <b>IHC:</b> 1:50 - 1:200 <b>IF/ICC:</b> 1:50 - 1:200
理论分子量	140kDa
实测分子量	140kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.75% BSA,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	293T
细胞定位	cytoplasm,cytosol,mitotic spindle,spindle pole
纯化	Affinity purification

## 抗原信息


抗原信息	Recombinant fusion protein corresponding to Human TACC3.
序列	MSLQVLNDKNVSNEKNTENCDFLSPPEVTGRSSVLRVSQKENVPPKNLAKAMKVTFQTPLRDPQTHRILSPSMASKLEA PFTQDDTLGLENSHPVWTQK

## 靶点信息


研究背景	This gene encodes a member of the transforming acidic coiled-coil protein family. The encoded protein is a motor spindle protein that may play a role in stabilization of the mitotic spindle. This protein may also play a role in growth and differentiation of certain cancer cells. [provided by RefSeq, Nov 2011]
基因ID	10460

基因名	TACC3
Swiss	Q9Y6A5
别名	ERIC-1; ERIC1

## 产品验证



Immunohistochemical analysis of paraffin-embedded human liver cancer, using TACC3 Antibody.



Western blot analysis of TACC3 expressed in 293T using TACC3 Rabbit mAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

## 实验步骤

访问官网浏览详情: [www.ablybio.cn](http://www.ablybio.cn)