

— ABLYBIO, Help Your Research



Atg4A Rabbit mAb

货号: **AYM31174**

产品信息

| | |
|-------|---|
| 反应 | Human,Mouse,Rat |
| 宿主 | Rabbit |
| 克隆性 | Monoclonal |
| 预测反应 | |
| 应用 | WB IHC IP |
| 推荐浓度 | WB: 1:500 - 1:2000 IHC: 1:50 - 1:200 IP: 1:20 - 1:50 |
| 理论分子量 | 36kDa/38kDa/42kDa/45kDa |
| 实测分子量 | 45kDa |
| 形式 | Liquid |
| 保存条件 | Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.75% BSA,50% glycerol,pH7.3. |
| 偶联物 | Unconjugated |
| 阳性对照 | HeLa,Mouse liver,Rat lung |
| 细胞定位 | Cytoplasm |
| 纯化 | Affinity purification |


抗原信息

| | |
|------|--|
| 抗原信息 | Recombinant fusion protein corresponding to Human Atg4A. |
|------|--|

靶点信息

| | |
|-------|--|
| 研究背景 | Autophagy is the process by which endogenous proteins and damaged organelles are destroyed intracellularly. Autophagy is postulated to be essential for cell homeostasis and cell remodeling during differentiation, metamorphosis, non-apoptotic cell death, and aging. Reduced levels of autophagy have been described in some malignant tumors, and a role for autophagy in controlling the unregulated cell growth linked to cancer has been proposed. This gene encodes a member of the autophagin protein family. The encoded protein is also designated as a member of the C-54 family of cysteine proteases. |
| 基因ID | 115201 |
| 基因名 | ATG4A |
| Swiss | Q8WYN0 |
| 别名 | Atg4A,Atg4A Rabbit mAb,AUT-like 2 cysteine endopeptidase,Autophagy-related cysteine endopeptidase 2,Autophagy-related protein 4 homolog A,APG4A |

产品验证



Western blot analysis of Atg4A expressed in HeLa, Mouse liver, Rat lung using Atg4A Rabbit mAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn (<https://www.ablybio.cn/>www.ablybio.cn)