

PCK1 Rabbit mAb

货号: **AYM31186**

产品信息

反应	Human
宿主	Rabbit
克隆性	Monoclonal
预测反应	
应用	WB IHC
推荐浓度	WB: 1:500 - 1:2000 IHC: 1:50 - 1:200
理论分子量	34kDa/69kDa
实测分子量	69kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.75% BSA,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	HepG2,Mouse liver,Mouse kidney,Rat liver,Rat pancreas,Rat kidney
细胞定位	Cytoplasm
纯化	Affinity purification

抗原信息


抗原信息	Recombinant fusion protein corresponding to Human PCK1.
序列	FGKYLAWLSMAQHAAKLPKIFHVNWFRKDKEGKFLWPGFGENSrvLEWmfNRIDGKASTKLTPIGYIPKEDALNLKGLGHINMMELFSISKEFWEKEVEDIEKYLEQVNADLPCEIEREILALKQRISQM

靶点信息

研究背景	This gene is a main control point for the regulation of gluconeogenesis. The cytosolic enzyme encoded by this gene, along with GTP, catalyzes the formation of phosphoenolpyruvate from oxaloacetate, with the release of carbon dioxide and GDP. The expression of this gene can be regulated by insulin, glucocorticoids, glucagon, cAMP, and diet. Defects in this gene are a cause of cytosolic phosphoenolpyruvate carboxykinase deficiency. A mitochondrial isozyme of the encoded protein also has been characterized.
基因ID	5105

基因名	PCK1
Swiss	P35558
别名	PCK1;PEPCK-C;PEPCK1;PEPCKC

产品验证



Western blot analysis of PCK1 expressed in HepG2, Mouse liver, Mouse kidney using PCK1 Rabbit mAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane . Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn