

Presenilin 1 Rabbit mAb

货号: **AYM31203**

产品信息

反应	Human,Mouse,Rat
宿主	Rabbit
克隆性	Monoclonal
预测反应	
应用	WB FC
推荐浓度	WB: 1:500 - 1:2000 FC: 1:20 - 1:50
理论分子量	21kDa/42kDa/46kDa/48kDa/52kDa
实测分子量	28kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.75% BSA,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	A-431,Jurkat,U-251MG,Mouse testis,Mouse liver,Mouse brain,Rat brain
细胞定位	Cell membrane,Cell surface,Endoplasmic reticulum membrane,Golgi apparatus membrane,Multi-pass membrane protein
纯化	Affinity purification


抗原信息

抗原信息	Recombinant fusion protein corresponding to Human Presenilin 1.
序列	MTELPAPLSYFQNAQMSEDNHLNNTVRSQNDNRERQEHNDRRSLGHPEPLSNGRPQNSRQVVEQDEEEDLTKYGAKHVIMLFVPVTLCMVVVVATI

靶点信息

研究背景	Alzheimer's disease (AD) patients with an inherited form of the disease carry mutations in the presenilin proteins (PSEN1; PSEN2) or in the amyloid precursor protein (APP). These disease-linked mutations result in increased production of the longer form of amyloid-beta (main component of amyloid deposits found in AD brains). Presenilins are postulated to regulate APP processing through their effects on gamma-secretase, an enzyme that cleaves APP. Also, it is thought that the presenilins are involved in the cleavage of the Notch receptor, such that they either directly regulate gamma-secretase activity or themselves are protease enzymes. Several alternatively spliced transcript variants encoding different isoforms have been identified for this gene, the full-length nature of only some have been determined. [provided by RefSeq, Aug 2008]
基因ID	5663
基因名	PSEN1
Swiss	P49768
别名	AD3;FAD;PS-1;PS1;S182;Presenilin 1;PSEN1

产品验证



Western blot analysis of Presenilin 1 expressed in Mouse testis, Rat brain, A-431 using Presenilin 1 Rabbit mAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn