— ABLYBIO, Help Your Research

# Nck Rabbit mAb

**货号: AYM31215**

## 产品信息

| | |
|---|---|
| 反应 | Human,Mouse,Rat |
| 宿主 | Rabbit |
| 克隆性 | Monoclonal |
| 预测反应 | |
| 应用 | WB IHC ICC IP FC |
| 推荐浓度 | **WB:** 1:500 - 1:2000<br>**IHC:** 1:50 - 1:200<br>**ICC:** 1:50 - 1:200<br>**IP:** 1:20 - 1:50<br>**FC:** 1:20 - 1:50 |
| 理论分子量 | 35kDa/42kDa |
| 实测分子量 | 47kDa |
| 形式 | Liquid |
| 保存条件 | Store at -20℃. Avoid freeze / thaw cycles.<br>Buffer: PBS with 0.75% BSA,50% glycerol,pH7.3. |
| 偶联物 | Unconjugated |
| 阳性对照 | Mouse brain |
| 细胞定位 | Cytoplasm,Endoplasmic reticulum,Nucleus |
| 纯化 | Affinity purification |


## 抗原信息

| | |
|---|---|
| 抗原信息 | Recombinant fusion protein corresponding to Human Nck. |

## 靶点信息

| | |
|---|---|
| 研究背景 | The protein encoded by this gene is one of the signaling and transforming proteins containing Src homology 2 and 3 (SH2 and SH3) domains. It is located in the cytoplasm and is an adaptor protein involved in transducing signals from receptor tyrosine kinases to downstream signal recipients such as RAS. Alternatively spliced transcript variants encoding different isoforms have been found. |
| 基因ID | 4690 |
| 基因名 | NCK1 |
| Swiss | P16333 (https://www.uniprot.org/uniprotkb/P16333/entry) |
| 别名 | Nck,Nck Rabbit mAb,NCK1,Cytoplasmic protein NCK1,NCK adapter protein 1,SH2/SH3 adapter protein NCK-alpha |

## 产品验证




Western blot analysis of Nck expressed in Mouse brain using Nck Rabbit mAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

## 实验步骤

访问官网浏览详情: www.ablybio.cn (https://www.ablybio.cn//www.ablybio.cn)