

Nck Rabbit mAb

货号: **AYM31215**

产品信息

反应	Human,Mouse,Rat
宿主	Rabbit
克隆性	Monoclonal
预测反应	
应用	WB IHC ICC IP FC
推荐浓度	WB: 1:500 - 1:2000 IHC: 1:50 - 1:200 ICC: 1:50 - 1:200 IP: 1:20 - 1:50 FC: 1:20 - 1:50
理论分子量	35kDa/42kDa
实测分子量	47kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.75% BSA,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	Mouse brain
细胞定位	Cytoplasm,Endoplasmic reticulum,Nucleus
纯化	Affinity purification


抗原信息

抗原信息	Recombinant fusion protein corresponding to Human Nck.
序列	RKNSARKASIVKNLKDTLGIGKVKRKPSVPDSASPADDSFVDPGERLYDLNMPAYVKFNMAEREDELKSLIKGTVIVMEK CSDGWWRGSGYNGQVGVFSPSNYVTEEGDSPLGDHVGSLSEKLAAVVNNLNTGQVLHVQALYPFSSSNDEELNFEKGD VMDVIEKPENDEWVKCRKINGMVGKLVKPNYVVMQNNPLTSGLESPPPQCDYIRPSLTGKFAGNPWYYGKVTRHQAE MALNERGHEGDFLIRDSESSPNDFSVSLKAQGNKHKFKVQLKETVYICIGQRKFSTMEELVEHYKKAPIFTSEQGEKLYLVK HLS

靶点信息

研究背景	The protein encoded by this gene is one of the signaling and transforming proteins containing Src homology 2 and 3 (SH2 and SH3) domains. It is located in the cytoplasm and is an adaptor protein involved in transducing signals from receptor tyrosine kinases to downstream signal recipients such as RAS. Alternatively spliced transcript variants encoding different isoforms have been found.
基因ID	4690
基因名	NCK1
Swiss	P16333
别名	NCK1;NCK;NCKalpha;nck-1

产品验证



Western blot analysis of Nck expressed in Mouse brain using Nck Rabbit mAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn