

CRKL Rabbit mAb

货号: **AYM31221**

产品信息

反应	Human
宿主	Rabbit
克隆性	Monoclonal
预测反应	
应用	WB IHC IF/ICC
推荐浓度	WB: 1:500 - 1:2000 IHC: 1:50 - 1:200 IF/ICC: 1:50 - 1:200
理论分子量	33kDa
实测分子量	39kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.75% BSA,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	LO2,Jurkat,SKOV3,Mouse brain,Rat heart
细胞定位	cytosol,extrinsic component of postsynaptic membrane,neuromuscular junction,nucleoplasm
纯化	Affinity purification

抗原信息


抗原信息	Recombinant fusion protein corresponding to Human CRKL.
序列	MSSARFDSSDRSAWYMGVSRQEAQTRLQGQRHGMFLVRDSSTCPGDYVLSVSENSRVSHYIINSLPNRRFKIGDQEFD HLPALLEFYKIHLYLDTTTLIEPAPRYSPPMGVSAPNLPTAEDNLEYVRTLYDFPGNDAEDLPFKKGEILVIIKPEEQWWSA RNKDGRVGMIPVYVEKLVRSPPHGKHGHRNSNSYGIPEPAHAYAQPQTTTLPVAVSGSPGAAITPLPSTQNGPVFAKAIQ KRVPCAYDKTALALEVGDIVKVTRMNINGQWEGEVNQRKGLFPFTHVKIFDPQNPDENE

靶点信息

研究背景	This gene encodes a protein kinase containing SH2 and SH3 (src homology) domains which has been shown to activate the RAS and JUN kinase signaling pathways and transform fibroblasts in a RAS-dependent fashion. It is a substrate of the BCR-ABL tyrosine kinase, plays a role in fibroblast transformation by BCR-ABL, and may be oncogenic.
------	---

基因ID	1399
基因名	CRKL
Swiss	P46109
别名	CRKL

产品验证



Western blot analysis of CRKL expressed in LO2, Jurkat, SKOV3 using CRKL Rabbit mAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn