

# STF1 Rabbit mAb

货号: **AYM31237**

## 产品信息

反应	Human,Mouse,Rat
宿主	Rabbit
克隆性	Monoclonal
预测反应	
应用	WB
推荐浓度	<b>WB:</b> 1:500 - 1:2000
理论分子量	52kDa
实测分子量	52kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.75% BSA,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	BxPC-3,Mouse ovary,Mouse spleen
细胞定位	Nucleus
纯化	Affinity purification

## 抗原信息


抗原信息	Recombinant fusion protein corresponding to Human STF1.
序列	AQELVLQLLALQLDRQEFVCLKFIILFSLDLKFLNNHILVKDAQEKANAALLDYTLCHYPHCGDKFQQLLLCLVEVRALSMQA KEYLYHKHLGNEMPRNLLIEMQLQAKQT

## 靶点信息

研究背景	The protein encoded by this gene is a transcriptional activator involved in sex determination. The encoded protein binds DNA as a monomer. Defects in this gene are a cause of XY sex reversal with or without adrenal failure as well as adrenocortical insufficiency without ovarian defect.
基因ID	2516
基因名	NR5A1

Swiss	Q13285
别名	NR5A1; AD4BP; ELP; FTZ1; FTZF1; POF7; SF-1; SF1; SPGF8; SRXY3; hSF-1; SRXX4; steroidogenic factor 1

## 产品验证



Western blot analysis of STF1 expressed in BxPC-3, Mouse ovary, Mouse spleen using STF1 Rabbit mAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane . Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

## 实验步骤

访问官网浏览详情: [www.ablybio.cn](http://www.ablybio.cn)